



SHAQUILLE GETS REAL ABOUT HEART FAILURE



A Guide for
African Americans
With **Heart Failure**

BiDil[®]
isosorbide dinitrate 20 mg/hydralazine HCl 37.5 mg

What Is Heart Failure (HF)?

HF doesn't mean your heart stops beating. It means it is not working very well. That causes problems pumping blood through your body. Over time, people with HF may feel worn out or short of breath. They may also have swelling in the feet and legs.¹

HF can sneak up on you. It doesn't happen all at once. At first, your heart tries to work harder. But it can't keep up over time. Once HF symptoms start, you need to see your doctor right away. Otherwise, they may get worse.

Shaq
BIDIL ACCESS
PROGRAM

Team Up With Shaq

Although I don't have heart failure, the issue hits close to home in my community. That's why I'm teaming up with Arbor to drive awareness about the first heart failure medication indicated for self-identified African American patients.

ShaqGetsReal.com

How HF Impacts the African American Community

HF is on the rise across America, including in the African American population.^{2,4} Risk factors for HF such as high blood pressure and diabetes occur at high rates in the African American community.^{2,3,5}

African Americans may be younger and sicker when they find out they have HF than whites⁶⁻¹⁰

Research suggests that African American patients are up to 20x more likely to get HF before age 50 than white Americans⁶

African Americans are hospitalized for HF more often than whites^{3,9}

Information for Patients about BiDil[®] (isosorbide dinitrate/hydralazine HCl)

BiDil is approved for use with other heart medicines to treat heart failure in black patients to improve survival, improve heart failure symptoms, and help patients stay out of the hospital longer. There is little experience in patients with heart failure who experience significant symptoms while at rest. Most patients in the clinical study of BiDil also received other heart failure medicines.

IMPORTANT SAFETY INFORMATION

Tell your doctor about any allergies you have, especially if you're sensitive to nitrates, such as nitroglycerin tablets or isosorbide dinitrate (Isordil[®]). BiDil[®] has a nitrate component, so you need to let your doctor know.

Tell your doctor if you're taking any erectile dysfunction or pulmonary hypertension drugs like Viagra[®] or Revatio[™] (sildenafil), Levitra[®] (vardenafil) or Cialis[®] (tadalafil).

Please see additional Important Safety Information and full Prescribing Information attached.

Ignoring HF Won't Make It Go Away

HF only gets worse with time. If you have shortness of breath or always feel tired, don't ignore it. HF needs to be treated quickly to get the best results. It is important to talk to your doctor and follow your heart failure treatment plan.¹

A HF Medicine Indicated Just for You

There is a medicine that is approved by FDA that can be added to other HF medicines in self-identified African Americans. That medicine is called BiDil[®] (isosorbide dinitrate/hydralazine HCl). BiDil was tested in a large clinical study that only included self-identified African American HF patients. The results showed that BiDil may help patients in this population live longer, feel better and delay first hospitalization for HF.^{11,12}

How BiDil Works

African American patients may have trouble with blood vessel relaxation.¹³ This makes it hard for blood to go through. Think of your blood vessel like a small pipe that's trying to get a lot of water through it. Researchers think that the 2 medicines in BiDil (isosorbide dinitrate/hydralazine HCl) help the blood vessels relax. This makes them wider and lets blood flow more easily.^{13,14}

The mechanism of action underlying the beneficial effects of BiDil in the treatment of HF has not been fully established.¹²

WARNINGS AND PRECAUTIONS


- Also tell your doctor if you are taking any medication to decrease blood pressure because when taken with BiDil, blood pressure may become too low.
- It is possible you'll get headaches, especially at first, but they often lessen over time. Keep your doctor posted on your headache progress; he or she may want to adjust your dosage.
- If you experience dizziness, call your doctor. Please make sure to tell your doctor about any of the signs or symptoms mentioned below or about any unusual events that worry you.

Please see additional Important Safety Information and full Prescribing Information attached.

TIGHTER BLOOD VESSEL



RELAXED BLOOD VESSEL



BiDil[®]
isosorbide dinitrate 20 mg/hydralazine HCl 37.5 mg

BiDil® Study Results

HF patients who took BiDil during the African-American Heart Failure Trial (A-HeFT) had:¹²

- 43% lower risk of dying
- 39% lower risk of going to the hospital for the first time for HF
- Significant improvement in their ability to function on a day-to-day basis (as reported by patients)

The main goal of the trial was to combine the above three pieces of information (death, first hospitalization for HF and day-to-day function) to see if BiDil made a difference. When these three items were combined, it showed BiDil made a significant difference.



A-HeFT was a double-blind study of 1050 self-identified African American patients that tested the efficacy of BiDil as an added therapy. In addition to their standard HF therapy, half of A-HeFT patients took BiDil and the other half took a placebo. As a double-blind study, neither the patients nor the prescribing physicians knew who was taking BiDil or the placebo.

The results were significant. In fact, BiDil reduced the rate of mortality so much vs the placebo, the study was concluded early to allow all study patients the opportunity to take BiDil.



Common Side Effects

Some people who take BiDil (isosorbide dinitrate/hydralazine HCl) get headaches at first. If headaches are still a problem, the doctor sometimes adjusts the BiDil dose. Other people have said they feel lightheaded or dizzy while on BiDil. This usually happens when they stand up after lying or sitting down. Headache and dizziness were the most common side effects of BiDil in clinical studies.¹²

Keep Taking Your Medicine

There's no cure for HF. So once you are on a HF medicine, make sure you keep taking it as your doctor directs, even if you are feeling better. Don't stop or change anything unless your doctor tells you to. Talk to your doctor before you stop taking it.¹⁵

WARNINGS AND PRECAUTIONS (cont'd)

- If you experience dizziness, call your doctor. Please make sure to tell your doctor about any of the signs or symptoms mentioned below or about any unusual events that worry you.
- Drinking less fluids than your doctor recommends or losing fluid due to diarrhea, sweating, or vomiting may cause low blood pressure, lightheadedness, or fainting. If fainting occurs, stop taking BiDil and contact your doctor immediately.
- Lightheadedness may occur when standing, especially after sitting or lying down.
- If you experience any achy and/or swollen joints, unexplained fever for more than a few days, skin rashes, chest pain, prolonged weakness or fatigue (even after a good night's sleep), or any other unexplained signs or symptoms, make sure to tell your doctor as they may be signs of a serious medical condition.
- You may also experience rapid heartbeat that could lead to chest pain or aggravate chest pain, or numbness or tingling in the hands or feet.

Please see additional Important Safety Information and full Prescribing Information attached.



Shaq BIDIL ACCESS PROGRAM

Together, let's learn more about managing HF and how it may be more affordable than ever.

ShaqGetsReal.com

BiDil® Patient Savings Card

Pay as little as \$XX for your first fill of BiDil (up to 90 tablets) and as little as \$XX for refills 2 through 12.

VISIT BIDIL.COM TO ENROLL

Medicare/Medicaid patients not eligible.

Preferred Arbor E-Z Rx Pharmacies

Pay just \$XX with insurance or \$XX cash for each monthly supply of BiDil (up to 90 tablets per month). Copay requirements apply.

**FIND PARTICIPATING PHARMACIES
AT EZRXLOCATOR.COM**

Medicare/Medicaid patients not eligible.

Cash Savings & Home Delivery

Arbor Patient Direct is a free premium prescription home delivery service that may be able to help patients willing to pay cash with prescription affordability. Pay just \$XX plus shipping for 90 tablets.

CALL 844-289-3981 TO ENROLL

Talking With Your Doctor About HF

When you visit the doctor, make sure you ask lots of questions. Don't hold back because you feel your question might be "silly." You need to know everything you can about HF so you can stay on track. And the best place to get that information is at the doctor.

Some things to talk with your doctor about might include:

How far along is my HF?

What problems will my HF cause going forward?

What caused my HF, and how are we fixing that?

What changes can I make to my daily life to help myself?

What are the best ways to make the changes I need to make?

What physical activities should I stay away from?

What kinds of physical activities can I still do?

Is there a way to make taking medicine on time easier?

What side effects from my medicines should I watch for?

What should I do if I feel worse all of a sudden?

Information for Patients About BiDil®

BiDil (isosorbide dinitrate/hydralazine HCl) is approved for use with other heart medicines to treat heart failure in black patients to improve survival, improve heart failure symptoms, and help patients stay out of the hospital longer. There is little experience in patients with heart failure who experience significant symptoms while at rest. Patients in the clinical study of BiDil received other heart failure medicines.

IMPORTANT SAFETY INFORMATION

Tell your doctor about any allergies you have, especially if you're sensitive to nitrates, such as nitroglycerin tablets or isosorbide dinitrate (Isordil®). BiDil has a nitrate component, so you need to let your doctor know.

Tell your doctor if you're taking any erectile dysfunction or pulmonary hypertension drugs like Viagra® or Revatio™ (sildenafil), Levitra® (vardenafil) or Cialis® (tadalafil).

WARNINGS AND PRECAUTIONS

Also tell your doctor if you are taking any medication to decrease blood pressure because when taken with BiDil, blood pressure may become too low.

It is possible you'll get headaches, especially at first, but they often lessen over time. Keep your doctor posted on your headache progress; he or she may want to adjust your dosage.

If you experience dizziness, call your doctor. Please make sure to tell your doctor about any of the signs or symptoms mentioned below or about any unusual events that worry you.

Drinking less fluids than your doctor recommends or losing fluid due to diarrhea, sweating, or vomiting may cause low blood pressure, lightheadedness, or fainting. If fainting occurs, stop taking BiDil and contact your doctor immediately.

Lightheadedness may occur when standing, especially after sitting or lying down.

If you experience any achy and/or swollen joints, unexplained fever for more than a few days, skin rashes, chest pain, prolonged weakness or fatigue (even after a good night's sleep), or any other unexplained signs or symptoms, make sure to tell your doctor as they may be signs of a serious medical condition.

You may also experience rapid heartbeat that could lead to chest pain or aggravate chest pain, or numbness or tingling in the hands or feet.

COMMON SIDE EFFECTS

Headache and dizziness were the most frequent side effects experienced in studies with BiDil.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information attached. This information does not take the place of talking with your healthcare provider about your condition or your treatment. Ask your doctor if BiDil may be right for you.

References: 1. American Heart Association. What is heart failure? http://www.heart.org/idc/groups/heart-public/@wcm/@hcm/documents/downloadable/ucm_300315.pdf. Published 2015. Accessed January 19, 2017. 2. Mitchell JE, Ferdinand KC, Watson KE, et al. Treatment of heart failure in African Americans—a call to action. *J Natl Med Assoc.* 2011;103(2):86-98. 3. Benjamin EJ, Virani SS, Callaway CW, et al. Heart disease and stroke statistics-2018 update: a report from the American Heart Association. *Circulation.* 2018;137(12):e67-e492. 4. Heidenreich PA, Albert NM, Allen LA, et al. Forecasting the impact of heart failure in the United States: A policy statement from the American Heart Association. *Circ Heart Fail.* May 2013;6(3):606-619. doi:1161/HHF.0b013e318291329a. 5. Yancy CW. Heart failure in African Americans: unique etiology and pharmacologic treatment responses. *J Natl Med Assoc.* 2003;95(1):1-9. 6. Bibbins-Domingo K, Pletcher MJ, Lin F, et al. Racial differences in incident heart failure among young adults. *N Engl J Med.* 2009;360(12):1179-1190. 7. Loefer LR, Rosamond WD, Chang PP, Folsom AR, Chambless LE. Heart failure incidence and survival (from the Atherosclerosis Risk in Communities study). *Am J Cardiol.* April 2008;101:1016-1022. doi:10.1016/j.amjcard.2007.11.061. 8. Bahrami H, Kronmal R, Bluemke DA, et al. Differences in the incidence of congestive heart failure by ethnicity: the multi-ethnic study of atherosclerosis. *Arch Intern Med.* October 2008;168(19):2138-2145. doi:10.1001/archinte.168.19.2138. 9. Dries DL, Exner DV, Gersh BJ, Cooper HA, Carson PE, Domanski MJ. Racial differences in the outcome of left ventricular dysfunction. *N Engl J Med.* February 1999;340(8):609-616. doi:10.1056/NEJM199902253400804. 10. Lee WC, Serag H, Ohsfeldt RL, et al. Racial Disparities in Type of Heart Failure and Hospitalization. *J Immigr Minor Health.* February 2019;21(1):98-104. doi:10.1007/s10903-018-0727-4. 11. Taylor AL, Ziesche S, Yancy C, et al. Combination of isosorbide dinitrate and hydralazine in blacks with heart failure. *New Engl J Med.* 2004;351(20):2049-2057. 12. BiDil [package insert]. Atlanta, GA: Arbor Pharmaceuticals, LLC. 13. Sharma A, Colvin-Adams M, Yancy CW. Heart failure in African Americans: disparities can be overcome. *Cleve Clin J Med.* 2014;81(5):301-311. 14. Yancy CW. Heart failure in African-Americans. *US Cardiol.* 2006;2005:2(1):196-200. 15. American Heart Association. Medications used to treat heart failure. http://www.heart.org/HEARTORG/Conditions/HeartFailure/TreatmentOptionsForHeartFailure/Medications-Used-to-Treat-Heart-Failure_UCM_306342_Article.jsp#.WC36jNyzgUI. Accessed November 17, 2016.

BiDil[®]

isosorbide dinitrate 20 mg/hydralazine HCl 37.5 mg

If You Are an African American With HF, Ask Your Doctor About BiDil[®] Today

- Proven to help self-identified African Americans with HF reduce risk of death and reduce risk of first hospitalization for HF
- Patients reported improvement in their day-to-day function according to a HF questionnaire
- Approved as an add-on to other HF medicines in self-identified African Americans
- Savings programs may help you get BiDil[®] (isosorbide dinitrate/hydralazine HCl) at a lower cost



SHAQUILLE GETS REAL ABOUT HEART FAILURE

ShaqGetsReal.com

COMMON SIDE EFFECTS

- Headache and dizziness were the most frequent side effects experienced in studies with BiDil.

IMPORTANT SAFETY INFORMATION

Tell your doctor about any allergies you have, especially if you're sensitive to nitrates, such as nitroglycerin tablets or isosorbide dinitrate (Isordil[®]). BiDil has a nitrate component, so you need to let your doctor know.

Tell your doctor if you're taking any erectile dysfunction or pulmonary hypertension drugs like Viagra[®] or Revatio[™] (sildenafil), Levitra[®] (vardenafil) or Cialis[®] (tadalafil).

Please see additional Important Safety Information and full Prescribing Information attached.

BiDil is a registered trademark of Arbor Pharmaceuticals, LLC. All other trademarks and registered trademarks are the property of their respective owners.


PHARMACEUTICALS, LLC

© 2019 Arbor Pharmaceuticals, LLC.
All rights reserved. PP-BDL-US-0690

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BDiDil safely and effectively. See full prescribing information for BDiDil.

BDiDil® (isosorbide dinitrate and hydralazine hydrochloride) tablet, for oral use

Initial U.S. Approval: 2005

INDICATIONS AND USAGE

BDiDil is a combination of isosorbide dinitrate, a nitrate vasodilator, and hydralazine hydrochloride, an arteriolar vasodilator, indicated for:

- the treatment of heart failure as an adjunct therapy to standard therapy in self-identified black patients to improve survival, prolong time to hospitalization for heart failure and to improve patient-reported functional status (1.1)

Limitations of use:

- There is little experience in patients with NYHA class IV heart failure (1.2)

DOSAGE AND ADMINISTRATION

One tablet three times a day titrated to a maximum tolerated dose up to two tablets three times a day (2)

Dosage may be decreased to as little as one-half tablet three times a day if intolerable side effects occur. Efforts should be made to titrate up as soon as side effects subside (2)

DOSAGE FORMS AND STRENGTHS

Tablets (scored): 20 mg isosorbide dinitrate and 37.5 mg hydralazine hydrochloride (3)

CONTRAINDICATIONS

- Patients who are allergic to organic nitrates (4)
- Use of phosphodiesterase type 5 (PDE5) inhibitors, such as avanafil, sildenafil, tadalafil, or vardenafil, or soluble guanylate cyclase (sGC) stimulator (riociguat). (4)

WARNINGS AND PRECAUTIONS

- May cause symptomatic hypotension (5.1)
- Symptomatic Lupus Erythematosus Syndromes:* Consider discontinuation if clinically appropriate (5.2)
- Myocardial ischemia and angina* (5.3)
- Peripheral Neuritis:* May be treated with Pyridoxine (5.4)

ADVERSE REACTIONS

Most common adverse reactions (≥ 5% more on BDiDil than on placebo) were headache and dizziness (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Arbor Pharmaceuticals, LLC at 1-866-516-4950 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- Geriatric Use:* Isosorbide dinitrate and hydralazine may be eliminated more slowly in elderly patients. Initiate therapy at the low end of the dosing range (8.5)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 03/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE**
 - Treatment of Heart Failure in Self-identified Black Patients
 - Limitations of Use
- DOSAGE AND ADMINISTRATION**
- DOSAGE FORMS AND STRENGTHS**
- CONTRAINDICATIONS**
- WARNINGS AND PRECAUTIONS**
 - Hypotension
 - Systemic Lupus Erythematosus
 - Worsening Ischemic Heart Disease
 - Peripheral Neuritis
- ADVERSE REACTIONS**
 - Clinical Trials Experience
 - Postmarketing Experience
- DRUG INTERACTIONS**
 - Phosphodiesterase Inhibitors
- USE IN SPECIFIC POPULATIONS**
 - Pregnancy
 - Lactation
 - Pediatric Use
 - Geriatric Use
 - Renal Impairment
 - Hepatic Impairment
- OVERDOSAGE**
- DESCRIPTION**
- CLINICAL PHARMACOLOGY**
 - Mechanism of Action
 - Pharmacodynamics
 - Pharmacokinetics
- NONCLINICAL TOXICOLOGY**
 - Carcinogenesis, Mutagenesis, Impairment of Fertility
- CLINICAL STUDIES**
- HOW SUPPLIED/STORAGE AND HANDLING**
- PATIENT COUNSELING INFORMATION**

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Treatment of Heart Failure in Self-identified Black Patients

BDiDil is indicated for the treatment of heart failure as an adjunct to standard therapy in self-identified black patients to improve survival, to prolong time to hospitalization for heart failure, and to improve patient-reported functional status.

1.2 Limitations of Use

There is little experience in patients with NYHA class IV heart failure.

2 DOSAGE AND ADMINISTRATION

BDiDil should be initiated at a dose of one BDiDil Tablet, three times a day. Titrate to a maximum of two tablets three times daily, if tolerated.

Although titration of BDiDil can be rapid (3-5 days), some patients may experience side effects and may take longer to reach their maximum tolerated dose. The dosage may be decreased to as little as one-half BDiDil Tablet three times a day if intolerable side effects occur. Efforts should be made to titrate up as soon as side effects subside.

3 DOSAGE FORMS AND STRENGTHS

The BDiDil (20 mg isosorbide dinitrate and 37.5 mg hydralazine hydrochloride) tablets are orange, biconvex, approximately 8 mm in diameter, scored, film-coated, and debossed with "20" on one side over the score and "N" on the other side.

4 CONTRAINDICATIONS

BDiDil is contraindicated in patients who are allergic to organic nitrates.

Do not use BDiDil in patients who are taking PDE-5 inhibitors, such as avanafil, sildenafil, tadalafil, or vardenafil. Concomitant use can cause severe hypotension, syncope, or myocardial ischemia [see *Drug Interactions* (7.1)].

Do not use BDiDil in patients who are taking the soluble guanylate cyclase (sGC) stimulator riociguat. Concomitant use can cause hypotension.

5 WARNINGS AND PRECAUTIONS

5.1 Hypotension

Symptomatic hypotension, particularly with upright posture, may occur with even small doses of BDiDil. Hypotension is most likely to occur in patients who have been volume or salt depleted; correct prior to initiation of BDiDil [see *Adverse Reactions* (6.1)].

5.2 Systemic Lupus Erythematosus

Hydralazine hydrochloride has been reported to cause a drug-induced systemic lupus erythematosus (SLE) syndrome. Symptoms and signs usually regress when hydralazine hydrochloride is discontinued.

5.3 Worsening Ischemic Heart Disease

Hydralazine hydrochloride can cause tachycardia and hypotension potentially leading to myocardial ischemia and angina, particularly in patients with hypertrophic cardiomyopathy.

5.4 Peripheral Neuritis

Hydralazine hydrochloride has been associated with peripheral neuritis, evidenced by paresthesia, numbness, and tingling, which may be related to an antipyridoxine effect. Pyridoxine should be added to BDiDil therapy if such symptoms develop.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

BDiDil has been evaluated for safety in 517 heart failure patients in A-HeFT. A total of 317 of these patients received BDiDil for at least 6 months, and 220 received BDiDil for at least 12 months. In A-HeFT, 21% of the patients discontinued BDiDil for adverse reactions compared to 12% who discontinued placebo. Overall, adverse reactions were more common in BDiDil-treated than in placebo-treated patients. Table 1 lists adverse reactions reported with an incidence, after rounding, ≥ 2% higher on BDiDil than on placebo in A-HeFT, regardless of causality. The most common reasons for discontinuing BDiDil in the A-HeFT trial was headache (7%).

Table 1. Adverse Reactions Occurring in the A-HeFT Study in ≥ 2% of Patients Treated with BDiDil.

	BiDil (N=517) %	Placebo (N=527) %
Headache	50	21
Dizziness	32	14
Asthenia	14	11
Nausea	10	6
Hypotension	8	4
Sinusitis	4	2
Ventricular tachycardia	4	2
Paresthesia	4	2
Vomiting	4	2
Amblyopia	3	1

In the V-HeFT I and II clinical studies, a total of 587 patients with heart failure were treated with the combination of isosorbide dinitrate and hydralazine hydrochloride. The type, pattern, frequency and severity of adverse reactions reported in these studies were similar to those reported in A-HeFT, described above and no unusual adverse reactions were reported.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of BDiDil. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Use of BDiDil: The following adverse reactions have been identified with use of BDiDil.

- Cardiac Disorders:* Palpitations
- Ear and labyrinth disorders:* Tinnitus, vertigo
- Eye Disorders:* Eyelid edema, vision blurred
- Gastrointestinal Disorders:* Abdominal discomfort, constipation
- General Disorders and Administration Site Conditions:* Facial pain, flushing, chest discomfort, chest pain, peripheral edema
- Musculoskeletal and Connective Tissue Disorders:* Pain in extremity, myalgia
- Nervous Disorders:* Dysgeusia, hypoesthesia, migraine, syncope
- Renal and Urinary Disorders:* Chromaturia, pulmonary renal syndrome
- Respiratory, Thoracic and Mediastinal Disorders:* Dyspnea
- Reproductive System and Breast Disorders:* Erectile dysfunction
- Skin and Subcutaneous Tissue Disorders:* Erythema, hyperhidrosis, pruritus, face swelling

Use of Hydralazine Hydrochloride or Isosorbide Dinitrate: The following reactions have been reported with use of either hydralazine hydrochloride or isosorbide dinitrate.

- Blood and Lymphatic System Disorders:* Blood dyscrasias, agranulocytosis, purpura, eosinophilia, splenomegaly.
- Eye Disorders:* Lacrimation, conjunctivitis.
- Gastrointestinal Disorders:* Paralytic ileus.
- Hepatobiliary Disorders:* Hepatitis.
- Psychiatric Disorders:* Psychotic reactions, disorientation.
- Renal and Urinary Disorders:* Difficulty in urination.

7 DRUG INTERACTIONS

7.1 Phosphodiesterase Inhibitors

BDiDil is contraindicated in patients who are using a selective inhibitor of cyclic guanosine monophosphate (cGMP)-specific phosphodiesterase type 5 (PDE5), PDE5 inhibitors such as avanafil, sildenafil, vardenafil, and tadalafil have been shown to potentiate the hypotensive effects of organic nitrates. Do not use BDiDil in patients who are taking the soluble guanylate cyclase (sGC) stimulator riociguat. Concomitant use can cause hypotension [see *Contraindications* (4)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no data on BDiDil use in pregnant women, and insufficient data on its components (hydralazine and isosorbide dinitrate) to assess a drug-associated risk of major birth defects or miscarriage with first trimester use. Available published data on hydralazine use in pregnancy during the second and third trimesters have not shown an association with adverse pregnancy-related outcomes.

Hydralazine hydrochloride is teratogenic in mice at 66 mg/kg and possibly in rabbits at 33 mg/kg (2 and 3 times the MRHD of BDiDil on a body surface area basis).

Isosorbide dinitrate has been shown to cause a dose-related increase in embryo-toxicity (excess mummified pups) in rabbits at 70 mg/kg (12 times the MRHD of BDiDil on a body surface area basis).

All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Clinical Considerations

Disease-Associated Maternal and/or Embryo/Fetal Risk

Pregnant women with heart failure are at increased risk for preterm birth. Clinical classification of heart disease may worsen with pregnancy and lead to maternal death and/or stillbirth.

8.2 Lactation

Risk Summary

There are no data on the presence of BDiDil in human or animal milk, the effects on the breastfed infant or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for BDiDil and any potential adverse effects on the breastfed child from BDiDil or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of BDiDil in children have not been established.

8.5 Geriatric Use

Clinical studies of BDiDil did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in response between elderly and younger patients. In general, dose selection for an elderly patient should start at the low end of the dosing range, reflecting the greater frequency of decreased hepatic and renal function, and of concomitant disease or other drug therapies.

Isosorbide dinitrate, its active metabolites, and hydralazine may be eliminated more slowly in elderly patients.

8.6 Renal Impairment

There are no studies of renal impairment using BDiDil. No dose adjustment is required for hydralazine or isosorbide dinitrate [see *Clinical Pharmacology* (12.3)].

Dialyzability of hydralazine has not been determined. Dialysis is not an effective method for removing isosorbide dinitrate or its metabolite isosorbide-5-mononitrate from the body.

8.7 Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of hydralazine alone has not been determined. Isosorbide dinitrate concentrations increase in patients with cirrhosis. There are no studies of hepatic impairment using BDiDil.

10 OVERDOSAGE

The signs and symptoms of overdosage with BDiDil are expected to be those of excessive pharmacologic effect, i.e., vasodilatation, reduced cardiac output and hypotension, and signs and symptoms include headache, confusion, tachycardia, and generalized skin flushing. Complications can include myocardial ischemia and subsequent myocardial infarction, cardiac arrhythmia, and profound shock. Syncope, coma and death may ensue without appropriate treatment.

Human Experience: There are no documented cases of overdosage with BDiDil. No deaths from acute poisoning have been reported.

Treatment: There is no specific antidote. Support of the cardiovascular system is of primary importance. Shock should be treated with plasma expanders, vasopressors, and positive inotropic agents. The gastric contents should be evacuated, taking adequate precautions to prevent aspiration. These manipulations have to be carried out after cardiovascular status has been stabilized, since they might precipitate cardiac arrhythmias or increase the depth of shock.

In patients with renal disease or congestive heart failure, therapy resulting in central volume expansion is not without hazard. Treatment of isosorbide dinitrate overdose in these patients may be difficult, and invasive monitoring may be required.


No data are available to suggest physiological maneuvers (e.g., maneuvers to change the pH of the urine) that might accelerate elimination of the components of BDiDil. Dialysis is not effective in removing circulating isosorbide dinitrate. The dialyzability of hydralazine has not been determined.

Methemoglobinemia: Nitrate ions liberated during metabolism of isosorbide dinitrate can oxidize hemoglobin into methemoglobin. There are case reports of significant methemoglobinemia in association with moderate overdoses of organic nitrates. Methemoglobin levels are measurable by most clinical laboratories. Methemoglobinemia could be serious in chronic heart failure patients because of already compromised vascular bed-tissue gas exchange dynamics. Classically, methemoglobinemic blood is described as chocolate brown, without color change on exposure to air. When methemoglobinemia is diagnosed, the treatment of choice is methylene blue, 1 to 2 mg/kg intravenously.

11 DESCRIPTION


BDiDil is a fixed-dose combination of isosorbide dinitrate, a vasodilator with effects on both arteries and veins, and hydralazine hydrochloride, a predominantly arterial vasodilator.

Isosorbide dinitrate is described chemically as 1,4,3,6-dianhydro-D-glucitol dinitrate and its structural formula is:



Isosorbide dinitrate is a white to off-white, crystalline powder with the empirical formula C₆H₈N₂O₈ and a molecular weight of 236.14. It is freely soluble in organic solvents such as alcohol, chloroform and ether, but is only sparingly soluble in water.

Hydralazine hydrochloride is described chemically as 1-hydrazinophthalazine monohydrochloride, and its structural formula is:



Hydralazine hydrochloride is a white to off-white, crystalline powder with the empirical formula C₈H₈N₄·HCl and a molecular weight of 196.64. It is soluble in water, slightly soluble in alcohol, and very slightly soluble in ether.

Each BDiDil Tablet for oral administration contains 20 mg of isosorbide dinitrate and 37.5 mg of hydralazine hydrochloride.

The inactive ingredients in BDiDil tablets include: anhydrous lactose, microcrystalline cellulose, sodium starch glycolate, colloidal silicon dioxide, magnesium stearate, hypromellose, FD&C Yellow No. 6 aluminum lake, polyethylene glycol, titanium dioxide, polysorbate 80.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action underlying the beneficial effects of BDiDil in the treatment of heart failure has not been established.

Isosorbide dinitrate is a vasodilator affecting both arteries and veins. Its dilator properties result from the release of nitric oxide and the subsequent activation of guanylyl cyclase, and ultimate relaxation of vascular smooth muscle.

Several well-controlled clinical trials have used exercise testing to assess the anti-anginal efficacy of chronically-delivered nitrates. In the large majority of these trials, active agents were no more effective than placebo after 24 hours (or less) of continuous therapy. Attempts to overcome nitrate tolerance by dose escalation, even to doses far in excess of those used acutely, have consistently failed. Only after nitrates have been absent from the body for several hours is response to nitrates restored.

Hydralazine hydrochloride is a selective dilator of arterial smooth muscle. Animal data suggests that hydralazine may also mitigate tolerance to nitrates.

12.2 Pharmacodynamics


The basis for the beneficial clinical effects of BDiDil is not known. In a small study of patients with chronic heart failure administered single doses of hydralazine 75 mg, isosorbide dinitrate 20 mg, and the combination, the combination elicited a statistically significant decrease in pulmonary capillary wedge pressure compared to hydralazine alone. The increase in cardiac output, renal blood flow and limb blood flow with the combination, however, was not greater than with hydralazine alone. There is no study of hemodynamic effects following multiple dosing.

12.3 Pharmacokinetics

Absorption

BiDil: Following a single 75-mg oral dose of hydralazine plus 40 mg of isosorbide dinitrate to 19 healthy adults, peak plasma concentrations of hydralazine (88 ng/mL/65 kg) and isosorbide dinitrate (76 ng/mL/65 kg) were reached in 1 hour. The half-lives were about 4 hours for hydralazine and about 2 hours for isosorbide dinitrate. Peak plasma concentrations of the two active metabolites, isosorbide-2-mononitrate and isosorbide-5-mononitrate, were 98 and 364 ng/mL/65 kg, respectively, at about 2 hours. No information is currently available regarding the effect of food on the bioavailability of hydralazine or isosorbide dinitrate from BDiDil tablets.

Hydralazine hydrochloride: About 2/3 of a 50-mg dose of ¹⁴C-hydralazine hydrochloride given in gelatin capsules was absorbed in hypertensive subjects. In patients with heart failure, mean absolute bioavailability of a single oral dose of hydralazine 75 mg varies from 10 to 26%, with the higher percentages in slow acetylators. Administration of doses escalating from 75 mg to 1000 mg three times daily to congestive heart failure patients resulted in an up to 9-fold increase in the dose normalized AUC, indicating non-linear kinetics of hydralazine, probably reflecting saturable first pass metabolism.



BDiDil®
(isosorbide dinitrate and hydralazine hydrochloride) Tablets

CIA75474F
BDL-PI-04 Rev. 03/2019

Isosorbide dinitrate: Absorption of isosorbide dinitrate from tablets after oral dosing is nearly complete. The average bioavailability of isosorbide dinitrate is about 25%, but is highly variable (10%-90%) because of first-pass metabolism, and increases progressively during chronic therapy. Serum concentrations reach their maximum about one hour after ingestion.

Distribution

Hydralazine hydrochloride: After intravenous administration of hydralazine in a dose of 0.3 mg/kg, the steady-state volume of distribution in patients with congestive heart failure was 2.2 L/kg.

Isosorbide dinitrate: The volume of distribution of isosorbide dinitrate is 2 to 4 L/kg. About 28% of circulating isosorbide dinitrate is protein bound.

Under steady-state conditions, isosorbide dinitrate accumulates significantly in muscle (pectoral) and vein (saphenous) wall relative to simultaneous plasma concentrations.

Metabolism

Hydralazine is metabolized by acetylation, ring oxidation and conjugation with endogenous compounds including pyruvic acid. Acetylation occurs predominantly during the first-pass after oral administration which explains the dependence of the absolute bioavailability on the acetylator phenotype. About 50% of patients are fast acetylators and have lower exposure.

After oral administration of hydralazine, the major circulating metabolites are hydralazine pyruvate hydrazone and methyltriazolophthalazine. Hydralazine is the main pharmacologically active entity; hydralazine pyruvate hydrazone has only minimal hypotensive and tachycardic activity. The pharmacological activity of methyltriazolophthalazine has not been determined. The major identified metabolite of hydralazine excreted in urine is acetylhydrazinophthalazine.

Isosorbide dinitrate undergoes extensive first-pass metabolism in the liver and is cleared at a rate of 2 to 4 L/minute with a serum half-life of about 1 hour. Isosorbide dinitrate's clearance is primarily by denitration to the 2-mononitrate (15 to 25%) and the 5-mononitrate (75 to 85%). Both metabolites have biological activity, especially the 5-mononitrate which has an overall half-life of about 5 hours. The 5-mononitrate is cleared by denitration to isosorbide, glucuronidation to the 5-mononitrate glucuronides, and by denitration/hydration to sorbitol. The 2-mononitrate appears to participate in the same metabolic pathways with a half-life of about 2 hours.

Elimination

Hydralazine: Metabolism is the main route for the elimination of hydralazine. Negligible amounts of unchanged hydralazine are excreted in urine.

Isosorbide dinitrate: Most isosorbide dinitrate is eliminated renally as conjugated metabolites.

Specific Populations

No pharmacokinetic studies in special populations were conducted with BiDil. Pharmacokinetics in special populations is based on individual components.

Geriatric Patients - The pharmacokinetics of hydralazine and isosorbide dinitrate, alone or in combination, have not been determined in patients over 65 years of age.

Pediatric Patients - The pharmacokinetics of hydralazine and isosorbide dinitrate, alone or in combination, have not been determined in patients below the age of 18 years.

Gender - There are no studies of gender-dependent effects with hydralazine. In a single dose study with isosorbide dinitrate, no gender-dependent differences in the pharmacokinetics of isosorbide dinitrate and its mononitrate metabolites were found.

Renal Impairment - The effect of renal impairment on the pharmacokinetics of hydralazine has not been determined. In a study with 49 hypertensive patients on chronic therapy with hydralazine in daily doses of 25-200 mg, the daily dose of hydralazine in 19 subjects with severely impaired renal function (creatinine clearance 5-28 mL/min) and in 17 subjects with normal renal function (creatinine clearance >100 mL/min) using a population PK approach was not different, suggesting no need for dose adjustment in patients with renal impairment. The dialyzability of hydralazine has not been determined. In three studies, renal insufficiency did not affect the pharmacokinetics of isosorbide dinitrate. Dialysis is not an effective method for removing isosorbide dinitrate or its metabolite isosorbide-5-mononitrate from the body.

Hepatic Impairment - The effect of hepatic impairment on the pharmacokinetics of hydralazine alone has not been determined. Isosorbide dinitrate concentrations increase in patients with cirrhosis.

Drug-Drug Interactions

No pharmacokinetic drug-drug interaction studies were conducted with BiDil.

Hydralazine: Administration of hydralazine can increase the exposure to a number of drugs including beta-blockers. In healthy males administered a single oral dose of hydralazine 50 mg and propranolol 1 mg/kg, the C_{max} and AUC for propranolol approximately doubled. In healthy subjects administered a single oral dose of hydralazine 50 mg and metoprolol 100 mg, the C_{max} and AUC for metoprolol increased by 50% and 30%, respectively. In pre-eclamptic women, twice-daily doses of hydralazine 25 mg and metoprolol 50 mg increased the C_{max} and AUC for metoprolol by 90% and 40%, respectively.

In healthy males administered single oral doses of hydralazine 25 mg and either lisinopril 20 mg or enalapril 20 mg, C_{max} and AUC for lisinopril were each increased 30%, but enalapril concentrations were unaffected.

Intravenous co-administration of 0.2 mg/kg hydralazine HCl and 40 mg furosemide in Japanese patients with congestive heart failure resulted in a 20% increase in the clearance of furosemide.

Isosorbide dinitrate: The vasodilating effects of coadministered isosorbide dinitrate may be additive to those of other vasodilators, including alcohol.

A single dose of 20 mg of isosorbide dinitrate was administered to healthy subjects after pretreatment with 80 mg propranolol three times daily for 48 hours, resulting in no impact on the pharmacokinetics of isosorbide dinitrate and isosorbide-5-mononitrate.

When single 100-mg oral doses of atenolol were administered 2 hours before isosorbide dinitrate at a 10-mg dose no differences in the pharmacokinetics of isosorbide dinitrate or its mononitrates were observed.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Hydralazine hydrochloride: An increased incidence of lung tumors (adenomas and adenocarcinomas) was observed in a lifetime study in Swiss albino mice given hydralazine hydrochloride continuously in their drinking water at a dosage of about 250 mg/kg per day (6 times the MRHD provided by BiDil on a body surface area basis). In a 2-year carcinogenicity study of rats given hydralazine hydrochloride by gavage at dose levels of 15, 30, and 60 mg/kg/day (up to 3 times the MRHD of BiDil on a body surface area basis), microscopic examination of the liver revealed a small, but statistically significant increase in benign neoplastic nodules in males (high-dosage) and females (both high and intermediate dosage groups). Benign interstitial cell tumors of the testes were also significantly increased in the high-dose group.

Hydralazine hydrochloride is mutagenic in bacterial systems, and is positive in rat and rabbit hepatocyte DNA repair studies *in vitro*. Additional *in vivo* and *in vitro* studies using lymphoma cells, germinal cells, fibroblasts from mice, bone marrow cells from Chinese hamsters and fibroblasts from human cell lines did not demonstrate any mutagenic or clastogenic potential for hydralazine hydrochloride.

Isosorbide dinitrate: No long-term animal studies have been performed to evaluate the mutagenic or carcinogenic potential of isosorbide dinitrate. A modified two-litter reproduction study among rats fed isosorbide dinitrate at 25 or 100 mg/kg/day (up to 9 times the Maximum Recommended Human Dose of BiDil on a body surface area basis) revealed no evidence of altered fertility or gestation.

14 CLINICAL STUDIES

BiDil or a combination of isosorbide dinitrate and hydralazine hydrochloride was studied in two placebo-controlled clinical trials in 1,692 patients with mild to severe heart failure (mostly NYHA class II and III) and one active control trial (vs. enalapril) in 804 patients. The results of the trials follow:

Placebo-controlled Study: In the multicenter trial V-HeFT I, the combination of hydralazine and isosorbide dinitrate 75 mg/40 mg 4 times daily (n=186) was compared to placebo (n=273) in men with impaired cardiac function and reduced exercise tolerance (primarily NYHA class II and III) and on therapy with digitalis glycosides and diuretics. There was no overall significant difference in mortality between the two treatment groups. There was, however, a trend favoring hydralazine and isosorbide dinitrate, which on retrospective analysis, was attributable to an effect in blacks (n=128). Survival in white patients (n=324) was similar on placebo and the combination treatment.

Active-controlled Study: In a second study of mortality, V-HeFT II, the combination of hydralazine and isosorbide dinitrate 75 mg/40 mg 4 times daily was compared to enalapril in 804 men with impaired cardiac function and reduced exercise tolerance (NYHA class II and III), and on therapy with digitalis glycosides and diuretics. The combination of hydralazine and isosorbide dinitrate was inferior to enalapril overall, but retrospective analysis showed that the difference was observed in the white population (n=574); there was essentially no difference in the black population (n=215).

Based on these retrospective analyses suggesting an effect on survival in black patients, but showing little evidence of an effect in the white population, a third study was conducted among black patients with heart failure.

Placebo-controlled Study: The A-HeFT trial evaluated BiDil vs. placebo among 1,050 self-identified black patients (over 95% NYHA class III) at 169 centers in the United States. All patients had stable symptomatic heart failure. Patients were required to have LVEF \leq 35% or left ventricular internal diastolic dimension $>$ 2.9 cm/m² plus LVEF $<$ 45%. Patients were maintained on stable background therapy and randomized to BiDil (n=518) or placebo (n=532). BiDil was initiated at 20 mg isosorbide dinitrate/37.5 mg hydralazine hydrochloride three times daily and titrated to a target dose of 40/75 mg three times daily or to the maximum tolerated dose. Patients were treated for up to 18 months.

The randomized population was 60% male, 1% NYHA class II, 95% NYHA class III and 4% NYHA class IV, with a mean age of 57 years, and was generally treated with standard treatments for heart failure including diuretics (94%, almost all loop diuretics), beta-blockers (87%), angiotensin converting enzyme inhibitors (ACE-I; 78%), angiotensin II receptor blockers (ARBs; 28%), either ACE-I or ARB (93%), digitalis glycosides (62%) and aldosterone antagonists (39%).

The primary endpoint was a composite score consisting of all-cause mortality, first hospitalization for heart failure, and responses to the Minnesota Living with Heart Failure questionnaire. The trial was terminated early, at a mean follow-up of 12 months, primarily because of a statistically significant 43% reduction in all-cause mortality in the BiDil-treated group (p=0.012; see Table 2 and Figure 1). The primary endpoint was also statistically in favor of BiDil (p \leq 0.021). The BiDil-treated group also showed a 39% reduction in the risk of a first hospitalization for heart failure (p<0.001; see Table 2 and Figure 2) and had statistically significant improvement in response to the Minnesota Living with Heart Failure questionnaire, a self-report of the patient's functional status, at most time points (see Figure 3). Patients in both treatment groups had mean baseline questionnaire scores of 51 (out of a possible 105).

Table 2. Results of A-HeFT (Intent-To-Treat Population)

	BiDil (N=518)	Placebo (N=532)	Hazard Ratio (95% CI)	P
Composite	-0.16 \pm 1.93	-0.47 \pm 2.04	—	0.021
All-cause mortality	6.2%	10.2%	0.57 (0.37, 0.89)	0.012
Hospitalization for heart failure	16.4%	24.4%	0.61 (0.46, 0.80)	<0.001

Figure 1: Kaplan-Meier Plot of Time to Death by All Cause in Black Patients (A-HeFT)




Figure 2: Kaplan-Meier Plot of Time to First Hospitalization for Heart Failure in Black Patients (A-HeFT)






Figure 3: Change in Minnesota Living With Heart Failure Score



Effects on survival and hospitalization for heart failure were similar in subgroups by age, gender, baseline disease, and use of concomitant medications, as shown in Figure 4.

Figure 4: Results for Demographic, Baseline Medication and Clinical Characteristic Subgroups in Black Patients (A-HeFT)



Patients treated with BiDil in the A-HeFT study had randomly measured blood pressures on average 3/3 mmHg lower than did patients on placebo. The contribution of the difference in blood pressure to the overall outcome difference is unknown. Whether both hydralazine and isosorbide dinitrate contribute to the overall outcome difference has not been studied in outcome trials. Isosorbide dinitrate and hydralazine have not been systematically studied for the treatment of heart failure as separate agents, and neither drug is indicated for heart failure.

16 HOW SUPPLIED/STORAGE AND HANDLING

BiDil Tablets contain 20 mg of isosorbide dinitrate and 37.5 mg of hydralazine hydrochloride. They are biconvex, approximately 8 mm in diameter, scored, film-coated, orange tablets debossed "20" on one side over the score and "N" on the other side.

- NDC 24338-010-18: Bottles of 180

Store at 25°C (77°F), excursions permitted to 15-30°C (59-86°F). [See USP Controlled Room Temperature.] Keep bottles tightly closed.

Protect from light. Dispense in a light-resistant, tight container.

17 PATIENT COUNSELING INFORMATION

Patients should be informed of possible side effects and advised to take the medication regularly and continuously as directed.

Headache

Inform patients that headaches often accompany treatment with BiDil, especially during initiation of treatment. Advise patients to consult a physician to adjust the dose of BiDil if headache continues with repeated dosing.

Hypotension

Warn patients about lightheadedness on standing.

Advise patients that inadequate fluid intake or excessive fluid loss from perspiration, diarrhea or vomiting may lead to an excessive fall in blood pressure and cause lightheadedness or even syncope. If syncope does occur, advise patients to discontinue BiDil and notify their prescribing physician as soon as possible.

Phosphodiesterase-5 Inhibitors

Advise patients to inform their physicians if they are taking, or planning to take, sildenafil, vardenafil, or tadalafil. BiDil should not be taken concomitantly with phosphodiesterase-5 inhibitors.

Worsening Ischemic Heart Disease

Advise patients to inform their physicians of any worsening of symptoms of myocardial ischemia, especially those with hypertrophic cardiomyopathy.

Systemic Lupus Erythematosus-like Symptoms

Advise patients if symptoms suggestive of systemic lupus erythematosus—such as arthralgia, fever, chest pain, prolonged malaise—occur to notify their prescribing physician.

Peripheral Neuritis

Advise patients if symptoms of peripheral neuritis—paresthesia, numbness, and tingling—occur to notify the prescribing physician.

Manufactured by:

Arbor
PHARMACEUTICALS, LLC
Atlanta, GA 30328

Manufactured by:

Lannett Company, Inc.
Philadelphia, PA 19136

BiDil is a registered trademark of

Arbor Pharmaceuticals, LLC

© 2019

U.S. Patent number 6,784,177; 6,465,463

All rights reserved

BDL-PI-04

CIA75474F