


Emgality is a preventive treatment specifically developed for migraine

Historically, many migraine preventive medicines were designed for other conditions. Emgality was specifically developed to bind to CGRP, a substance in the brain that may play a key role in migraine.

CGRP=calcitonin gene-related peptide.



Select Safety Information

Emgality may cause allergic reactions, such as itching, rash, hives, and trouble breathing. Allergic reactions can happen days after using Emgality. Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms, which may be part of an allergic reaction:

- swelling of your face, mouth, tongue, or throat
- trouble breathing

Please see Purpose and Safety Summary on page 8. See Instructions for Use included with the device.

Experience Emgality for as little as \$0 for up to 12 months with the Emgality Savings Card*



NO THE PHARMACES

These cells must be accompanied by a valid prescription for Empilety (galaneseumin-grini) and care only be used by the second must be accompanied by the second property of the second prope

*Terms and Conditions apply.

Visit Emgality.com/savings to download and activate the Savings Card or call 1-833-EMGALITY (1-833-364-2548) or text "info" to 55900. Msg & data rates may apply.

Once activated, present your Savings Card to the pharmacist when you pick up your prescription.

Please see the full Terms and Conditions <u>here</u>.



If you have additional questions, call your doctor or call the Emgality Answers Center at 1-833-EMGALITY (1-833-364-2548), Monday through Friday from 8 AM to 7 PM ET.

Emgality helps prevent future migraine days, so you can make the most of them

As shown in two 6-month studies of adults who had 4-14 migraine days per month:

Emgality CUT MIGRAINE DAYS IN HALF or more

or about 60% of people vs about 39% who took placebo

In a 3-month study of adults with 15+ headache days per month. 28% had their monthly migraine days cut in half or more with Emgality compared to 15% who took placebo.

- pregnant or plan to become pregnant. It is not known if Emgality will harm your unborn baby
- taking any prescription or over-the-counter medicines,

Please see Purpose and Safety Summary on page 8.

MIGRAINE-FREE MONTH

12% and 16% on average vs 6% who took placebo

In a 3-month study, the number of adults with 15+ headache days per month who were migraine-free in an average month did not differ between Emgality and placebo.

Select Safety Information

Before you use Emgality, tell your healthcare provider if you are:

- breastfeeding or plan to breastfeed. It is not known if Emgality passes into breast milk
- vitamins, or herbal supplements

See Instructions for Use included with the device.

Once-monthly Emgality comes in an easy-to-use, self-injectable Pen that can be used in the comfort of your home





The first time you inject Emgality, you will receive 2 Pens that should be administered together. After that, it's just 1 Pen, once a month.



In a study, 94% of people agreed at first use that the Emgality Pen was easy to use, and 96% were confident in their ability to use the Pen.





Receive the injection how-to video: Text INFO to 55900 to get the video delivered straight to your phone. Msg & data rates may apply.

Select Safety Information

The most common side effects of Emgality include injection site reactions. Tell your healthcare provider if you have any side effect that bothers you or does not go away. These are not all the possible side effects of Emgality.



3 basic steps to inject with the Emgality Pen

Before you inject for the first time, your healthcare provider should show you how to prepare and inject Emgality.







2. Place and unlock



3. Press, hold, and listen

IMPORTANT: These are not the complete instructions for using your device. Please see Instructions for Use that come with your device or visit Emgality.com. The most common side effects of Emgality are injection site reactions such as pain, redness, and itching.

After using your Emgality Pen, dispose of it safely in a sharps container. **Do not** throw the Emgality Pen in the trash. If you do not have a puncture-resistant container with a secure lid, order a free one at **Emgality.com/taking-emgality**.

Please see Purpose and Safety Summary on page 8. See Instructions for Use included with the device.

Have questions? We have answers



Get help when you need it

If you have additional questions, call your doctor or the Emgality Answers Center at 1-833-EMGALITY (1-833-364-2548), Monday through Friday from 8 AM to 7 PM ET. Our healthcare professionals are ready to walk you through the injection process and answer any questions you may have as you get started on Emgality.

Text to receive the injection how-to video

Text INFO to 55900 to get the injection how-to video delivered straight to your phone.

Msg & data rates may apply.

Share your story

Your story can help others with migraine get the answers they need. Start sharing by calling us at **1-833-EMGALITY (1-833-364-2548)**.



PURPOSE and SAFETY SUMMARY

Important Facts About Emgality® (em-GAL-it-ē) injection. Also known as galcanezumab-gnlm.

Emgality is a prescription medicine used for the preventive treatment of migraine in adults. The medicine comes in a prefilled pen or syringe and is taken once a month.

Warnings

Do not use Emgality if you are allergic to galcanezumab-gnlm or any of the ingredients in Emgality.

Emgality may cause allergic reactions, such as itching, rash, hives, and trouble breathing. Allergic reactions can happen days after using Emgality. Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms, which may be part of an allergic reaction: swelling of your face, mouth, tongue, or throat, or trouble breathing.

Common side effects

The most common side effects of Emgality are injection site reactions.

These are not all the possible side effects of Emgality. Tell your doctor if you have any side effects.

You can report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.

Before using

Before you use Emgality, discuss these questions with your doctor:

- ☐ Do you have other medical conditions?
- ☐ Do you take other medicines? Tell your doctor about all the prescription and over-the-counter drugs, vitamins, and herbal supplements you take.

Tell your doctor if:

- ☐ You are pregnant or breastfeeding.
- ☐ You are under age 18. It is not known if Emgality is safe and effective in children.

How to take

- Read the Instructions for Use that come with Emgality. There you will find information about how to store, prepare, and inject Emgality.
- Use Emgality exactly as your doctor says.
- If you miss a dose, inject it as soon as possible.
 Then continue to take it once a month from the date of your last dose.

Learn more

For more information, go to www.emgality.com or call 1-833-EMGALITY (1-833-364-2548) Monday - Friday, 8 AM - 7 PM EST.

This summary provides basic information about Emgality, but it does not include all information known about this medicine. Read the information that comes with your prescription every time you have a prescription filled. Talk to your doctor or pharmacist about this information. Your doctor is the best person to help you decide if Emgality is right for you.

GZ CON MIG BS 19FEB2020

Please see Instructions for Use that come with the device.



do with vour

MIGRAINE-FREE

DAYS?





Emgality* is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates. PP-GZ-US-1238 04/2020 ©Lilly USA, LLC 2020. All rights reserved.